



Product Information

Phaseolus vulgaris lectin E (PHA-E), Red kidney bean, Peroxidase conjugate

Cat. No.: XLC-56-14Q-E

Size: 1 mg

Synonym: PHA-E; Red-kidney-bean-agglutinin

This product is for research use only and is not intended for diagnostic use.

Product Information

Description	<i>Phaseolus vulgaris</i> lectin E (PHA-E) specially binds to Galactose and Complex structures. Inhibiting or Eluting Sugar: Bovine thyroglobulin or Acetic acid. This product can be used in glycobiology, immunohistochemistry, immunoassay and western blot.
Biological Source	Red kidney bean
Concentration	1 mg/mL
Conjugate	Peroxidase
Carbohydrate Specificity	Galactose; Complex structures
Inhibiting or Eluting Sugar	Bovine thyroglobulin or Acetic acid
Divalent Ions	Ca ⁺⁺ , Mn ⁺⁺
Form	Liquid
Purity	Pure
Potency	PHA-E (< 5 µg/mL) agglutinates type O human erythrocytes; PHA-E (< 0.5 mg/mL) agglutinates neuraminidase treated erythrocytes.
Buffer	0.01 M PBS (pH 7.2)
Preservative	No preservatives
Applications	Glycobiology, Immunohistochemistry, Immunoassay, Western blot
Storage	Store at -20°C. Protect from light.