



Product Information

Triticum vulgare lectin (WGA), Wheat germ, Acrylamide Beads conjugate

Cat. No.: XLC-71-19Q

Size: 2 mL

Synonym: WGA; Wheat germ agglutinin

This product is for research use only and is not intended for diagnostic use.

Product Information

Description	<i>Triticum vulgare</i> lectin (WGA) specially binds to <i>N</i> -Acetylglucosamine. Inhibiting or Eluting Sugar: Chitin hydrolysate or <i>N</i> -Acetylglucosamine w/acid or salt. This product can be used in glycobiology, immunochemistry, western blot and enzyme-linked immunosorbent assay.
Biological Source	Wheat germ
Concentration	2-2.5 mg/mL
Conjugate	Acrylamide Beads
Carbohydrate Specificity	<i>N</i> -Acetylglucosamine
Inhibiting or Eluting Sugar	Chitin hydrolysate or <i>N</i> -Acetylglucosamine w/acid or salt
Blood Group Specificity	Non-specific
Form	Liquid
Buffer	0.01 M PBS (pH 7.2)
Preservative	0.05% sodium azide
Applications	Glycobiology, Immunochemistry, Western blot, Enzyme-linked immunosorbent assay
Storage	Store at 2-8°C. (Do not freeze!)